arXiv:1206.3510v1 [math.CO] 15 Jun 2012

AN EXPANSION FORMULA FOR THE INVERSIONS AND EXCEDANCES IN THE SYMMETRIC GROUP

JIANG ZENG

ABSTRACT. We prove a recent conjecture of Blanco and Petersen (arXiv:1206.0803v2) about an expansion formula for inversions and excedances in the symmetric group.

1. INTRODUCTION

Let $\mathfrak{S}_n$ be the set of permutations on $[n] = \{1, \dots, n\}$. For $\sigma = \sigma(1) \dots \sigma(n) \in \mathfrak{S}_n$, the entry $i \in [n]$ is called an *excedance* (resp. *drop*) of σ if $i < \sigma(i)$ (resp. $i > \sigma(i)$); the entry $i \in [n]$ is called a *descent* of σ if $i < n$ and $\sigma(i) > \sigma(i+1)$. Denote the number of descents, excedances and drops in σ by $\operatorname{des}\sigma$, $\operatorname{exc}\sigma$ and $\operatorname{drop}\sigma$, respectively. It is well known [FS70] that the statistics des, exc and drop have the same distribution on $\mathfrak{S}_n$, their common enumerative polynomial is called the Eulerian polynomial:

$$A_n(t) = \sum_{\sigma \in \mathfrak{S}_n} t^{\operatorname{des}\sigma} = \sum_{\sigma \in \mathfrak{S}_n} t^{\operatorname{exc}\sigma} = \sum_{\sigma \in \mathfrak{S}_n} t^{\operatorname{drop}\sigma}, \tag{1}$$

and there are nonnegative integers $\gamma_{n,j}$ such that

$$A_n(t) = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} \gamma_{n,k} t^k (1+t)^{n-1-2k}. \tag{2}$$

In the last years several q-analogues of (2) were established in [SW10, HJZ12, SZ12] by combining one of the Eulerian statistics in (1) and the Major index. Let $\operatorname{inv}\sigma$ be the number of inversions in $\sigma \in \mathfrak{S}_n$. The following q-Eulerian polynomial

$$S_n(q,t) = \sum_{\sigma \in \mathfrak{S}_n} q^{\operatorname{inv}\sigma} t^{\operatorname{exc}\sigma} \tag{3}$$

was first studied in [CSZ97]. Recently, Blanco and Petersen [BP12, Conj. 3.1] has conjectured that $S_n(q,t)$ would display a nice q-version of (2).

Conjecture 1 (Blanco and Petersen)**.** *There exist polynomials $\gamma_{n,k}(q)$ with nonnegative integer coefficients such that:*

$$S_n(q,t/q) := \sum_{\sigma \in \mathfrak{S}_n} q^{\operatorname{inv}\sigma - \operatorname{exc}\sigma} t^{\operatorname{exc}\sigma} = \sum_{0 \le k \le (n-1)/2} \gamma_{n,k}(q) t^k (1+t)^{n-1-2k}. \tag{4}$$

The aim of this paper is to show that a stronger version of the above conjecture was implicitly given in [SZ10, SZ12].

Date: March 8, 2013.

2. Prelimaries

For $\sigma \in \mathfrak{S}_n$, the statistic $(31\text{-}2)\,\sigma$ (resp. $(13\text{-}2)\,\sigma$) is the number of pairs (i,j) such that $2 \leq i < j \leq n$ and $\sigma(i-1) > \sigma(j) > \sigma(i)$ (resp. $\sigma(i-1) < \sigma(j) < \sigma(i)$). Similarly, the statistic $(2\text{-}13)\,\sigma$ (resp. $(2\text{-}31)\,\sigma$) is the number of pairs (i,j) such that $1 \leq i < j \leq n-1$ and $\sigma(j+1) > \sigma(i) > \sigma(j)$ (resp. $\sigma(j+1) < \sigma(i) < \sigma(j)$). Define the generalized Eulerian polynomial by

$$A_n(p,q,t) := \sum_{\sigma\in\mathfrak{S}_n} p^{(2\text{-}13)\,\sigma} q^{(31\text{-}2)\,\sigma} t^{\operatorname{des}\sigma}. \tag{5}$$

For $\sigma \in \mathfrak{S}_n$, the value $\sigma(i)$, for $i \in [n]$, is called

- a *valley* if $\sigma(i-1) > \sigma(i)$ and $\sigma(i) < \sigma(i+1)$;
- a *double descent* if $\sigma(i-1) > \sigma(i)$ and $\sigma(i) > \sigma(i+1)$.

where $\sigma(0) = \sigma(n+1) = 0$ by convention. Let $\mathfrak{S}_{n,k}$ be the subset of permutations $\sigma \in \mathfrak{S}_n$ with exactly k valleys and without double descents. Define the polynomial

$$a_{n,k}(p,q) = \sum_{\sigma\in\mathfrak{S}_{n,k}} p^{(2\text{-}13)\,\sigma} q^{(31\text{-}2)\,\sigma}. \tag{6}$$

The following lemma is a special case of Theorem 2 in [SZ12].

Lemma 2. *We have the expansion formula*

$$A_n(p,q,t) = \sum_{k=0}^{\lfloor (n-1)/2\rfloor} a_{n,k}(p,q) t^k (1+t)^{n-1-2k}. \tag{7}$$

Moreover, for all $0 \leq k \leq \lfloor (n-1)/2 \rfloor$*, the following divisibility holds*

$$(p+q)^k \mid a_{n,k}(p,q). \tag{8}$$

For a permutation $\sigma \in \mathfrak{S}_n$ the crossing and nesting numbers are defined by

$$\operatorname{cros}\sigma = \#\left\{(i,j)\in[n]\times[n] : (i<j\leq\sigma(i)<\sigma(j)) \vee (i>j>\sigma(i)>\sigma(j))\right\}, \tag{9}$$

$$\operatorname{nest}\sigma = \#\left\{(i,j)\in[n]\times[n] : (i<j\leq\sigma(j)<\sigma(i)) \vee (i>j>\sigma(j)>\sigma(i))\right\}. \tag{10}$$

For example, if the permutation is $\sigma = 3762154 = \left(\begin{smallmatrix} 1&2&3&4&5&6&7\\ 3&7&6&2&1&5&4\end{smallmatrix}\right)$, there are three crossings $(1 < 2 < \sigma(1) < \sigma(2))$, $(1 < 3 \leq \sigma(1) < \sigma(3))$, $(7 > 4 > \sigma(7) > \sigma(4))$ and three nestings $(2 < 3 < \sigma(3) < \sigma(2))$, $(5 > 4 > \sigma(4) > \sigma(5))$, $(7 > 6 > \sigma(6) > \sigma(7))$, thus $\operatorname{cros}\sigma = \operatorname{nest}\sigma = 3$.

The following lemma is a special case of Theorem 5 in [SZ12].

Lemma 3. *There is a bijection* Φ *on* $\mathfrak{S}_n$ *such that for all* $\sigma \in \mathfrak{S}_n$ *we have*

$$(\operatorname{nest}, \operatorname{cros}, \operatorname{drop})\sigma = (2\text{-}31, 31\text{-}2, \operatorname{des})\Phi(\sigma).$$

The following lemma is proved in [SZ12, Eq. (39)] using continued fractions of their generating functions.

Lemma 4. *The triple statistics* $((2\text{-}31), (31\text{-}2), \operatorname{des})$ *and* $((2\text{-}13), (31\text{-}2), \operatorname{des})$ *are equidistributed on* $\mathfrak{S}_n$.

3. Proof of the conjecture

We first state a more precise version of Conjecture 1.

Theorem 5. *The Conjecture 1 holds true. Moreover, for* $0 \leq k \leq (n-1)/2$,

$$\gamma_{n,k}(q) = a_{n,k}(q^2, q) = \sum_{\sigma\in\mathfrak{S}_{n,k}} q^{2\,(\text{2-13})\,\sigma + (\text{31-2})\,\sigma}$$

and $\gamma_{n,k}(q)$ *is divisible by* $q^k(1+q)^k$.

Proof. Since $\operatorname{inv}\sigma^{-1} = \operatorname{inv}\sigma$ and $\operatorname{drop}\sigma = \operatorname{exc}\sigma^{-1}$ for any $\sigma \in \mathfrak{S}_n$, we have

$$S_n(q,t) = \sum_{\sigma\in\mathfrak{S}_n} q^{\operatorname{inv}\sigma} t^{\operatorname{exc}\sigma} = \sum_{\sigma\in\mathfrak{S}_n} q^{\operatorname{inv}\sigma} t^{\operatorname{drop}\sigma}.$$

Combining Lemmas 3 and 4 yields

$$A_n(p,q,t) = \sum_{\sigma\in\mathfrak{S}_n} p^{\operatorname{nest}\sigma} q^{\operatorname{cros}\sigma} t^{\operatorname{drop}\sigma}.$$

Therefore, as $\operatorname{inv} = \operatorname{drop} + \operatorname{cros} + 2\operatorname{nest}$ (see [SZ10, Eq. (40)]), we deduce

$$S_n(q, t/q) = A_n(q^2, q, t).$$

The desired result follows then from Lemma 2. $\square$

Remark. We conclude this paper with the following open problems:

1) Is there a simple biective proof of Lemma 4?
2) Is there a type B analogue of (2) for inversions and excedances?

References

[BP12] S. Blanco and T. Kyle Petersen, Counting Dyck paths by area and rank, arXiv:1206.0803v2.

[CSZ97] R. J. Clarke, E. Steingrímsson, and J. Zeng, New Euler-Mahonian statistics on permutations and words, Adv. in Appl. Math. **18** (1997), no. 3, 237–270.

[FS70] D. Foata and M.-P. Schützenberger, *Théorie géométrique des polynômes eulériens*, Lecture Notes in Mathematics, Vol. 138, Springer-Verlag, Berlin, 1970.

[HJZ12] G.-N. Han, F. Jouhet, J. Zeng, Two new triangles of q-integers via q-Eulerian polynomials of type A and B, arXiv:1203.6736.

[SW10] J. Shareshian and M. L. Wachs, Eulerian quasisymmetric functions, Adv. in Math., 225 (2010), 2921–2966.

[SZ10] H. Shin and J. Zeng, The q-tangent and q-secant numbers via continued fractions, European J. Combin. **31** (2010), no. 7, 1689–1705.

[SZ12] ———, The symmetric and unimodal expansion of Eulerian polynomials via continued fractions. European J. Combin. 33 (2012), no. 2, 111–127.

(Jiang Zeng) Université de Lyon; UMR 5208 du CNRS; Université Lyon 1; Institut Camille Jordan; 43, blvd du 11 novembre 1918, F-69622 Villeurbanne Cedex, France

E-mail address: `zeng@math.univ-lyon1.fr`